



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

Display Summary Show: 20 Sort Send to Text

About Entrez

Items 1 - 20 of 208

Page 1 of 11 Next

Text Version

## Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

## PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

## Related Resources

Order Documents  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

☐ 1: [Selber K, Collen A, Hyytia T, Penttila M, Tjerneld F, Kula MR.](#) Related Articles, Links



Parameters influencing protein extraction for whole broths in detergent based aqueous two-phase systems.

Bioseparation. 2001;10(4-5):229-36.

PMID: 12233746 [PubMed - indexed for MEDLINE]

☐ 2: [Collen A, Selber K, Hyytia T, Persson J, Nakari-Setälä T, Bailey M, Fagerström R, Kula MR, Penttila M, Stalbrand H, Tjerneld F.](#) Related Articles, Links



Primary recovery of a genetically engineered *Trichoderma reesei* endoglucanase I (Cel 7B) fusion protein in cloud point extraction systems.

Biotechnol Bioeng. 2002 May 20;78(4):385-94.

PMID: 11948445 [PubMed - indexed for MEDLINE]

☐ 3: [Collen A, Persson J, Linder M, Nakari-Setälä T, Penttila M, Tjerneld F, Sivars U.](#) Related Articles, Links



A novel two-step extraction method with detergent/polymer systems for primary recovery of the fusion protein endoglucanase I-hydrophobin I.

Biochim Biophys Acta. 2002 Jan 15;1569(1-3):139-50.

PMID: 11853968 [PubMed - indexed for MEDLINE]

☐ 4: [Collen A, Penttila M, Stalbrand H, Tjerneld F, Veide A.](#) Related Articles, Links



Extraction of endoglucanase I (Cel 7B) fusion proteins from *Trichoderma reesei* culture filtrate in a poly(ethylene glycol)-phosphate aqueous two-phase system.

J Chromatogr A. 2002 Jan 11;943(1):55-62.

PMID: 11820281 [PubMed - indexed for MEDLINE]

☐ 5: [Selber K, Müller S, Gieren H, Thommes J, Sottmann T, Strey R, Kula MR.](#) Related Articles, Links



Physical properties of detergent-based aqueous two-phase systems.

Bioseparation. 2001;10(4-5):243-53.

PMID: 12233748 [PubMed - indexed for MEDLINE]

☐ 6: [Collen A, Ward M, Tjerneld F, Stalbrand H.](#) Related Articles, Links



Genetically engineered peptide fusions for improved protein partitioning in aqueous two-phase systems. Effect of fusion localization on endoglucanase I of *Trichoderma reesei*.

J Chromatogr A. 2001 Mar 2;910(2):275-84.

PMID: 11261722 [PubMed - indexed for MEDLINE]











☐ 7: [Collen A, Ward M, Tjerneld F, Stalbrand H.](#) Related Articles, Links



Genetic engineering of the *Trichoderma reesei* endoglucanase I (Cel 7B) for enhanced partitioning in aqueous two-phase systems containing thermoseparating ethylene oxide--propylene oxide copolymers.

J Biotechnol. 2001 May 4;87(2):179-91.

PMID: 11278040 [PubMed - indexed for MEDLINE]

- ☐ **8:** [Rodenbrock A, Selber K, Egmond MR, Kula MR.](#) Related Articles, Links  
 Extraction of peptide tagged cutinase in detergent-based aqueous two-phase systems.  
Bioseparation. 2000;9(5):269-76.  
PMID: 11394565 [PubMed - indexed for MEDLINE]
- ☐ **9:** [Terstappen GC, Ramelmeier RA, Kula MR.](#) Related Articles, Links  
 Protein partitioning in detergent-based aqueous two-phase systems.  
J Biotechnol. 1993 Apr;28(2-3):263-75.  
PMID: 7763561 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Ramelmeier RA, Terstappen GC, Kula MR.](#) Related Articles, Links  
 The partitioning of cholesterol oxidase in Triton X-114-based aqueous two-phase systems.  
Bioseparation. 1991;2(5):315-24.  
PMID: 1368208 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Linder M, Selber K, Nakari-Setälä T, Qiao M, Kula MR, Penttilä M.](#) Related Articles, Links  
 The hydrophobins HFBI and HFBII from *Trichoderma reesei* showing efficient interactions with nonionic surfactants in aqueous two-phase systems.  
Biomacromolecules. 2001 Summer;2(2):511-7.  
PMID: 11749214 [PubMed - indexed for MEDLINE]
- ☐ **12:** [Venancio A, Teixeira JA, Mota M.](#) Related Articles, Links  
 Evaluation of crude hydroxypropyl starch as a bioseparation aqueous-phase-forming polymer.  
Biotechnol Prog. 1993 Nov-Dec;9(6):635-9.  
PMID: 7764352 [PubMed - indexed for MEDLINE]
- ☐ **13:** [Juhasz T, Szengyel Z, Szijarto N, Reczey K.](#) Related Articles, Links  
 Effect of pH on cellulase production of *Trichoderma reesei* RUT C30.  
Appl Biochem Biotechnol. 2004 Spring;113-116:201-11.  
PMID: 15054207 [PubMed - indexed for MEDLINE]
- ☐ **14:** [Kepka C, Collet E, Persson J, Stahl A, Lagerstedt T, Tjerneld F, Veide A.](#) Related Articles, Links  
 Pilot-scale extraction of an intracellular recombinant cutinase from *E. coli* cell homogenate using a thermoseparating aqueous two-phase system.  
J Biotechnol. 2003 Jun 26;103(2):165-81.  
PMID: 12814875 [PubMed - indexed for MEDLINE]
- ☐ **15:** [Rito-Palomares M, Negrete A, Galindo E, Serrano-Carreón L.](#) Related Articles, Links  
 Aroma compounds recovery from mycelial cultures in aqueous two-phase processes.  
J Chromatogr B Biomed Sci Appl. 2000 Jun 23;743(1-2):403-8.  
PMID: 10942311 [PubMed - indexed for MEDLINE]
- ☐ **16:** [Costa MJ, Cunha MT, Cabral JM, Aires-Barros MR.](#) Related Articles, Links  
 Scale-up of recombinant cutinase recovery by whole broth extraction with PEG-phosphate aqueous two-phase.  
Bioseparation. 2000;9(4):231-8.  
PMID: 11321521 [PubMed - indexed for MEDLINE]
- ☐ **17:** [Selber K, Nellen F, Steffen B, Thommes J, Kula MR.](#) Related Articles, Links  
 Investigation of mathematical methods for efficient optimisation of aqueous two-phase extraction.

J Chromatogr B Biomed Sci Appl. 2000 Jun 23;743(1-2):21-30.  
PMID: 10942269 [PubMed - indexed for MEDLINE]

☐ 18: [Sivars U, Tjerneld F.](#)

[Related Articles, Links](#)



Mechanisms of phase behaviour and protein partitioning in detergent/polymer aqueous two-phase systems for purification of integral membrane proteins.  
Biochim Biophys Acta. 2000 Apr 6;1474(2):133-46.  
PMID: 10742592 [PubMed - indexed for MEDLINE]

☐ 19: [Fujita Y, Ito J, Ueda M, Fukuda H, Kondo A.](#)

[Related Articles, Links](#)



Synergistic saccharification, and direct fermentation to ethanol, of amorphous cellulose by use of an engineered yeast strain codisplaying three types of cellulolytic enzyme.  
Appl Environ Microbiol. 2004 Feb;70(2):1207-12.  
PMID: 14766607 [PubMed - indexed for MEDLINE]

☐ 20: [Sadana A.](#)

[Related Articles, Links](#)



Protein inactivations during novel bioseparation techniques.  
Bioseparation. 1994 Feb;4(1):39-61.  
PMID: 7764586 [PubMed - indexed for MEDLINE]

Items 1 - 20 of 208

Page 1 of 11 Next

Display Summary Show: 20 Sort Send to Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 23 2004 06:26:50



Entrez PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed for   

Limits

Preview/Index

History

Clipboard

Details

  Show:    

About Entrez

Items 21 - 40 of 208

Previous   of 11 Next

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ **21:** [Wohlfahrt G, Pellikka T, Boer H, Teeri TT, Koivula A.](#)[Related Articles, Links](#)**Probing pH-dependent functional elements in proteins: modification of carboxylic acid pairs in *Trichoderma reesei* cellobiohydrolase Cel6A.**

Biochemistry. 2003 Sep 2;42(34):10095-103.

PMID: 12939137 [PubMed - indexed for MEDLINE]

☐ **22:** [Diamond AD, Hsu JT.](#)[Related Articles, Links](#)**Aqueous two-phase systems for biomolecule separation.**

Adv Biochem Eng Biotechnol. 1992;47:89-135. Review.

PMID: 1456110 [PubMed - indexed for MEDLINE]

☐ **23:** [Cunha MT, Costa MJ, Calado CR, Fonseca LP, Aires-Barros MR, Cabral JM.](#)[Related Articles, Links](#)**Integration of production and aqueous two-phase systems extraction of extracellular *Fusarium solani* pisi cutinase fusion proteins.**

J Biotechnol. 2003 Jan 9;100(1):55-64.

PMID: 12413786 [PubMed - indexed for MEDLINE]

☐ **24:** [Brumbauer A, Johansson G, Reczey K.](#)[Related Articles, Links](#)**Fractionation of cellulase and beta-glucosidase in a *Trichoderma reesei* culture liquid by use of two-phase partitioning.**

Bioseparation. 1999;7(6):287-95.

PMID: 10643639 [PubMed - indexed for MEDLINE]

☐ **25:** [Fernandes S, Mattiasson B, Hatti-Kaul R.](#)[Related Articles, Links](#)**Recovery of recombinant cutinase using detergent foam.**

Biotechnol Prog. 2002 Jan-Feb;18(1):116-23.

PMID: 11822909 [PubMed - indexed for MEDLINE]

☐ **26:** [Linder MB, Qiao M, Laumen F, Selber K, Hyytia T, Nakari-Setälä T, Penttilä ME.](#) [Related Articles, Links](#)**Efficient purification of recombinant proteins using hydrophobins as tags in surfactant-based two-phase systems.**

Biochemistry. 2004 Sep 21;43(37):11873-82.

PMID: 15362873 [PubMed - indexed for MEDLINE]

☐ **27:** [Sanchez-Ferrer A, Bru R, Garcia-Carmona F.](#)[Related Articles, Links](#)**Phase separation of biomolecules in polyoxyethylene glycol nonionic detergents.**

Crit Rev Biochem Mol Biol. 1994;29(4):275-313. Review.

PMID: 8001397 [PubMed - indexed for MEDLINE]

☐ **28:** [Costa SA, Pessoa A Jr, Roberto IC.](#)[Related Articles, Links](#)**Partitioning of xylanolytic complex from *Penicillium janthinellum* by an aqueous two-phase system.**

J Chromatogr B Biomed Sci Appl. 2000 Jun 23;743(1-2):339-48.  
PMID: 10942304 [PubMed - indexed for MEDLINE]

- ☐ 29: [Linder M, Szilvay GR, Nakari-Setälä T, Söderlund H, Penttilä M.](#)

[Related Articles, Links](#)



Surface adhesion of fusion proteins containing the hydrophobins HFBI and HFBII from *Trichoderma reesei*.

Protein Sci. 2002 Sep;11(9):2257-66.

PMID: 12192081 [PubMed - indexed for MEDLINE]

- ☐ 30: [Boer H, Koivula A.](#)

[Related Articles, Links](#)



The relationship between thermal stability and pH optimum studied with wild-type and mutant *Trichoderma reesei* cellobiohydrolase Cel7A.

Eur J Biochem. 2003 Mar;270(5):841-8.

PMID: 12603317 [PubMed - indexed for MEDLINE]

- ☐ 31: [Gunduz U.](#)

[Related Articles, Links](#)



Optimization of bovine serum albumin partition coefficient in aqueous two-phase systems.

Bioseparation. 2000;9(5):277-81.

PMID: 11394566 [PubMed - indexed for MEDLINE]

- ☐ 32: [Persson J, Nystrom L, Ageland H, Tjerneld F.](#)

[Related Articles, Links](#)



Purification of recombinant apolipoprotein A-1 Milano expressed in *Escherichia coli* using aqueous two-phase extraction followed by temperature-induced phase separation.

J Chromatogr B Biomed Sci Appl. 1998 Jun 26;711(1-2):97-109.

PMID: 9699979 [PubMed - indexed for MEDLINE]

- ☐ 33: [Paloheimo M, Mantyla A, Kallio J, Suominen P.](#)

[Related Articles, Links](#)



High-yield production of a bacterial xylanase in the filamentous fungus *Trichoderma reesei* requires a carrier polypeptide with an intact domain structure.

Appl Environ Microbiol. 2003 Dec;69(12):7073-82.

PMID: 14660351 [PubMed - indexed for MEDLINE]

- ☐ 34: [Forney CE, Glatz CE.](#)

[Related Articles, Links](#)



Extraction of charged fusion proteins in reversed micelles: comparison between different surfactant systems.

Biotechnol Prog. 1995 May-Jun;11(3):260-4.

PMID: 7619396 [PubMed - indexed for MEDLINE]

- ☐ 35: [Abonyo BO, Wang P, Narasaraaju TA, Rowan WH 3rd, McMillan DH, Zimmerman UJ, Liu L.](#)

[Related Articles, Links](#)



Characterization of alpha-soluble N-ethylmaleimide-sensitive fusion attachment protein in alveolar type II cells: implications in lung surfactant secretion.

Am J Respir Cell Mol Biol. 2003 Sep;29(3 Pt 1):273-82. Epub 2003 Mar 27.

PMID: 12663329 [PubMed - indexed for MEDLINE]

- ☐ 36: [Gunduz U.](#)

[Related Articles, Links](#)



Partitioning of bovine serum albumin in an aqueous two-phase system: optimization of partition coefficient.


J Chromatogr B Biomed Sci Appl. 2000 Jun 23;743(1-2):259-62.

PMID: 10942295 [PubMed - indexed for MEDLINE]

- ☐ 37: [Kamei DT, King JA, Wang DI, Blankschtein D.](#)


[Related Articles, Links](#)

Understanding viral partitioning in two-phase aqueous nonionic micellar systems: 2.

-  **Effect of entrained micelle-poor domains.**  
Biotechnol Bioeng. 2002 Apr 20;78(2):203-16.  
PMID: 11870611 [PubMed - indexed for MEDLINE]


☐ **38:** [Tsao GT, Gong CS, Cao NJ.](#)

[Related Articles, Links](#)

-  **Repeated solid-phase fermentation and extraction for enzyme production.**  
Appl Biochem Biotechnol. 2000 Spring;84-86:505-24.  
PMID: 10849816 [PubMed - indexed for MEDLINE]


☐ **39:** [Hitscherich C Jr, Aseyev V, Wiencek J, Loll PJ.](#)

[Related Articles, Links](#)

-  **Effects of PEG on detergent micelles: implications for the crystallization of integral membrane proteins.**  
Acta Crystallogr D Biol Crystallogr. 2001 Jul;57(Pt 7):1020-9.  
PMID: 11418772 [PubMed - indexed for MEDLINE]

☐ **40:** [Forciniti D, Hall CK, Kula MR.](#)

[Related Articles, Links](#)

-  **Temperature dependence of the partition coefficient of proteins in aqueous two-phase systems.**  
Bioseparation. 1991;2(2):115-28.  
PMID: 1368076 [PubMed - indexed for MEDLINE]


Items 21 - 40 of 208

Previous **Page**  of 11 Next

**Display**  ☒ Show:  ☒ Sort  ☒ **Send to**  ☒

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 23 2004 06:26:50



Entrez PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed ☒ for hydrophobin AND isolation  ☒ Limits Preview/Index History Clipboard Details

Limits: Publication Date to 1999/03

 Summary ☒ Show: 20 ☒ Sort ☒ Send to Text ☒

Items 1 - 9 of 9

One page.

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: [Lugones LG, Wosten HA, Wessels JG.](#)[Related Articles, Links](#)

A hydrophobin (ABH3) specifically secreted by vegetatively growing hyphae of *Agaricus bisporus* (common white button mushroom).

Microbiology. 1998 Aug;144 ( Pt 8):2345-53.

PMID: 9720057 [PubMed - indexed for MEDLINE]

☐ 2: [Tenberge KB, Stellamanns P, Plenz G, Robenek H.](#)[Related Articles, Links](#)

Nonradioactive in situ hybridization for detection of hydrophobin mRNA in the phytopathogenic fungus *Claviceps purpurea* during infection of rye.

Eur J Cell Biol. 1998 Mar;75(3):265-72.

PMID: 9587058 [PubMed - indexed for MEDLINE]

☐ 3: [Spanu P.](#)[Related Articles, Links](#)

HCF-1, a hydrophobin from the tomato pathogen *Cladosporium fulvum*.

Gene. 1997 Jul 1;193(1):89-96.

PMID: 9249071 [PubMed - indexed for MEDLINE]

☐ 4: [Lugones LG, Bosscher JS, Scholtmeyer K, de Vries OM, Wessels JG.](#)[Related Articles, Links](#)

An abundant hydrophobin (ABH1) forms hydrophobic rodlet layers in *Agaricus bisporus* fruiting bodies.

Microbiology. 1996 May;142 ( Pt 5):1321-9.

PMID: 8704971 [PubMed - indexed for MEDLINE]

☐ 5: [De Groot PW, Schaap PJ, Sonnenberg AS, Visser J, Van Griensven LJ.](#)[Related Articles, Links](#)

The *Agaricus bisporus* hypA gene encodes a hydrophobin and specifically accumulates in peel tissue of mushroom caps during fruit body development.

J Mol Biol. 1996 Apr 19;257(5):1008-18.

PMID: 8632464 [PubMed - indexed for MEDLINE]

☐ 6: [Nakari-Setälä T, Aro N, Kalkkinen N, Alatalo E, Penttilä M.](#)[Related Articles, Links](#)

Genetic and biochemical characterization of the *Trichoderma reesei* hydrophobin HFBI.

Eur J Biochem. 1996 Jan 15;235(1-2):248-55.

PMID: 8631337 [PubMed - indexed for MEDLINE]

☐ 7: [Templeton MD, Greenwood DR, Beever RE.](#)[Related Articles, Links](#)

Solubilization of *Neurospora crassa* rodlet proteins and identification of the predominant protein as the proteolytically processed eas (ccg-2) gene product.

Exp Mycol. 1995 Jun;19(2):166-9.

PMID: 7614378 [PubMed - indexed for MEDLINE]

☐ 8: [Bidochka MJ, St Leger RJ, Joshi L, Roberts DW.](#)[Related Articles, Links](#)

An inner cell wall protein (cwp1) from conidia of the entomopathogenic fungus *Beauveria*



bassiana.

Microbiology. 1995 May;141 ( Pt 5):1075-80.

PMID: 7773402 [PubMed - indexed for MEDLINE]

☐ 9: [Wosten HA, Schuren FH, Wessels JG.](#)

[Related Articles, Links](#)



Interfacial self-assembly of a hydrophobin into an amphipathic protein membrane mediates fungal attachment to hydrophobic surfaces.

EMBO J. 1994 Dec 15;13(24):5848-54.

PMID: 7813424 [PubMed - indexed for MEDLINE]

**Display**  ☐ Show:  ☐ Sort  ☐ **Send to**  ☐

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 23 2004 06:26:50